

Masses of Black Holes in Active Galactic Nuclei

Bradley M. Peterson

Department of Astronomy, The Ohio State University, 140 West 18th Avenue, Columbus, OH 43210, USA

Abstract. We present a progress report on a project whose goal is to improve both the precision and accuracy of reverberation-based black-hole masses. Reverberation masses appear to be accurate to a factor of about three, and the black-hole mass/bulge velocity dispersion ($M-\sigma$) relationship appears to be the same in active and quiescent galaxies.

1. The Importance of Reverberation Masses

A detailed knowledge of the masses of supermassive black holes (SBHs) in AGNs is an important element of our understanding of how the accretion process works in nature. Highly accurate (to 10% or so) masses are needed to understand the energy budgets in AGNs: how much of the energy is converted into radiation, how much appears in the form of kinetic energy in jets and winds, and how much (if any) is advected in the black hole itself? Less accurate mass determinations (to within a factor of a few or so) are nevertheless very useful for current-epoch black-hole demographics that allow us to approach the same questions about accretion physics from a statistical point of view. Current estimates of black-hole masses are anchored by nearly 40 AGNs (Wandel, Peterson, & Malkan 1999; Kaspi et al. 2000; Onken et al. 2002) whose SBH masses have been estimated by emission-line reverberation methods (Blandford & McKee 1982; Peterson 1993, 2001), in which the response time of the broad emission lines to continuum variations is taken to be the light-travel time across the line-emitting region. By combining this length scale with a measurement of the line width, a virial mass estimate is obtained through

$$M = \frac{kc\tau V^2}{G}, \quad (1)$$

where c and G are the speed of light and gravitational constant, respectively, τ is the response time scale for a broad emission line, V is a suitable measure of the emission-line width, and k is a numerical factor that depends on the currently unknown structure and kinematics of the line-emitting region. There are two strong pieces of evidence that suggest that reverberation masses are reliable to within a factor of a few:

1. In four AGNs for which multiple emission-line lags and line widths have been measured, a virial relationship between lag and line width (i.e., $\tau \propto V^{-2}$) is seen (Peterson & Wandel 1999, 2000; Onken & Peterson 2002).

2. AGN SBH masses seem to follow the same relationship between black-hole mass and host galaxy bulge velocity dispersion, the M – σ relationship (Ferrarese & Merritt 2000; Gebhardt et al. 2000), that is seen in quiescent galaxies (Ferrarese et al. 2001, and references therein).

Masses obtained by reverberation methods are becoming increasingly important as they are used to anchor scaling methods that allow estimation of the black-hole masses in higher-redshift, higher-luminosity AGNs (e.g., Wandel, Peterson, & Malkan 1999; Vestergaard 2002; McLure & Jarvis 2002) which thus allows us in principle to probe the growth of quasar masses over most of the history of the Universe. Furthermore, unlike stellar-dynamical and gas-dynamical methods, the reverberation mapping technique is directly applicable to higher-redshift sources because it is independent of angular resolution.

Given the potential importance of the reverberation masses for these sources and the simple fact that a similar or superior sample is not likely to be obtained in the immediate future, we decided to undertake a detailed reanalysis of the existing reverberation data to improve on the homogeneity and accuracy of the database. In this contribution, we describe some preliminary results.

2. Refining Reverberation Masses

Our intent is to improve both the accuracy and precision to which reverberation masses are known. The specific goals of this investigation are:

1. *To minimize the formal errors in BLR radius and line width determinations.* The goal is to increase the measured *precision* of the virial product rV^2 . We assume that the virial relationship holds, and test various characterizations of the emission-line time lag and velocity width to minimize the scatter in the virial product for AGNs for which multiple measurements are available.
2. *To determine the best characterization of the black-hole mass.* The goal is to improve the *accuracy* of reverberation masses by determining the constant k in eq. (1). Here we assume that AGNs and quiescent galaxies follow the same M – σ relationship and that k can be determined by scaling the AGNs relative to quiescent black holes. In sacrificing the chance of finding systematic differences between active and quiescent galaxies, we are able to constrain the BLR geometry and kinematics in a statistical sense by determining both k and its corresponding uncertainty, i.e., the scatter in the AGN M – σ relationship.

3. Some Preliminary Results

As we will describe in detail elsewhere, we conducted tests based on both numerical simulations and different treatments of actual monitoring data. The large database on NGC 5548 (see Peterson et al. 2002 and references therein), for which there are 14 years of optical monitoring data plus two years of UV monitoring, plays a particularly important role in this analysis. We examined




Figure 1. The solid line is the M – σ relationship for quiescent galaxies, which are shown as small filled circles. The large triangles are AGNs from this work, with $k = 9.57$ (see eq. 1). The open triangles are NGC 7469, NGC 5548, and Akn 120 (left to right), which have the highest precision virial products and dominate the determination of k .

the data carefully to determine which measures of lag and line width give the most consistent virial product rV^2 . We reach the following conclusions:

1. In general, the centroid of the cross-correlation function is a more robust estimate of the lag than the peak value of the cross-correlation.
2. The width of an emission-line in the rms spectrum¹ better characterizes the line-of-sight velocity dispersion than does the line width in the mean spectrum.
3. The line dispersion (i.e., the second moment of the emission-line profile) is superior to the full-width at half maximum (FWHM) as a measure of the line-of-sight velocity width. However, both measures give a slope consistent with $\tau \propto V^{-2}$.

4. The M – σ Relationship for AGNs

If we make the reasonably well-justified assumption that AGNs follow the same M – σ relationship as quiescent galaxies, then we can compute the scale factor k in eq. (1) by scaling the raw virial products rV^2 to the nominal quiescent-galaxy M – σ relationship. In a statistical sense, this can provide us with an

¹The multiple spectra obtained in a reverberation-mapping experiment can be used to compute both a mean and root-mean-square spectrum. The latter isolates the variable part of the spectrum.

admittedly quite weak statistical constraint on the structure of the broad-line region. The result of such a scaling exercise is shown in Fig. 1, for which we find $k = 9.57 \pm 0.15$. This weighted fit is dominated by Akn 120, NGC 5548, and NGC 7469, the three AGNs with the highest-precision virial products. The scatter of the AGNs around the M – σ relationship is about 0.5 dex; i.e., reverberation masses appear to be accurate to a factor of about three.

For comparison, we note that the values of k for various trivial BLR models are $k = 3$ for an isothermal sphere or for a thin spherical shell of randomly inclined Keplerian orbits, and $k = 2/\sin^2 i$ for a ring at inclination i . In the latter case, a typical inclination of $i \approx 27^\circ$ would yield our present estimate of k . The fact that our inferred value of k is in the expected range for simple models reassures us that our assumption that the M – σ relationships for active and quiescent galaxies are identical is probably not grossly in error.

Acknowledgments. The work described here is being undertaken with my colleagues K.M. Gilbert, C.A. Onken, R.W. Pogge, and M. Vestergaard (Ohio State), L. Ferrarese and D. Merritt (Rutgers), S. Kaspi, D. Maoz, and H. Netzer (Tel-Aviv Univ.), and A. Wandel (Hebrew Univ.). We are grateful for funding of this work through NASA grant NAG5-8397, National Science Foundation grant AST-0205964, and US-Israel Binational Science Foundation grant 1999336.

References

- Blandford, R., & McKee, C.F. 1982, *ApJ*, 255, 419
- Ferrarese, L. 2002, in *Current High-Energy Emission Around Black Holes*, ed. C.-H. Lee (Singapore: World Scientific), in press (astro-ph/0203047)
- Ferrarese, L., & Merritt, D. 2000, *ApJ*, 539, L9
- Ferrarese, L., Pogge, R.W., Peterson, B.M., Merritt, D., Wandel, A., & Joseph, C.L. 2001, *ApJ*, 555, L79
- Gebhardt, K., et al. 2000, *ApJ*, 539, 13
- Kaspi, S., et al. 2000, *ApJ*, 533, 631
- McLure, R.J., & Jarvis, M.J. 2002, *MNRAS*, in press (astro-ph/0204473)
- Onken, C.A., & Peterson, B.M. 2002, *ApJ*, 572, 746
- Onken, C.A., Peterson, B.M., Dietrich, M., Robinson, A., & Salamanca, I. 2002, submitted to *ApJ*
- Peterson, B.M. 1993, *PASP*, 105, 247
- Peterson, B.M. 2001, in *Advanced Lectures on the Starburst–AGN Connection*, ed. I. Aretxaga, D. Kunth, & R. Mújica, (Singapore: World Scientific), 3
- Peterson, B.M., et al. 2002, *ApJ*, in press (astro-ph/0208064)
- Peterson, B.M., & Wandel, A. 1999, *ApJ*, 521, L95
- Peterson, B.M., & Wandel, A. 2000, *ApJ*, 540, L13
- Vestergaard, M. 2002, *ApJ*, 571, 733
- Wandel, A., Peterson, B.M., & Malkan, M.A. 1999, *ApJ*, 526, 579